# **PBA Taylor**<sup>(D)</sup> 'Kaspa-type' field pea

# Virus resistant kaspa-type



#### MAIN ADVANTAGES

PBA Taylor<sup>(b)</sup> (tested as OZP1408) is a mid flowering and early to mid maturing field pea, slightly later than PBA Wharton<sup>(b)</sup> but earlier than Kaspa<sup>(b)</sup>. It is the highest yielding field pea variety across most cropping regions as of 2021 and outclasses the existing dual virus resistant variety PBA Wharton<sup>(b)</sup> except in regions with high boron and salinity constraints.

PBA Taylor<sup>(b)</sup> has the same plant type as Kaspa<sup>(b)</sup> – semileafless and semi-dwarf plant architecture with nonshattering pods and Kaspa-type seed. PBA Taylor<sup>(b)</sup> has resistance to two virus diseases: pea seed borne mosaic virus (PSbMV) and bean leaf roll virus (BLRV).

### **SEED PROTECTION & ROYALTIES**

PBA Taylor<sup>(b)</sup> is protected under Plant Breeder's Rights (PBR) legislation. Growers can only retain seed from their production of PBA Taylor<sup>(b)</sup> for own seed use.

An End Point Royalty (EPR) of \$2.97 per tonne (GST inclusive), which includes breeder royalties, applies upon delivery of this variety. Seed is available from the commercial partner Seednet.




## **KEY FEATURES**

- Consistently highest yielding field pea variety across many environments.
- Mid flowering and early to mid maturity.
- Similar erect and semi-leafless plant type to Kaspa<sup>()</sup>.
- Resistant to pea seed-borne mosaic virus and bean leaf roll virus.
- Grain marketable as 'Kaspa-type'.

#### **AREA OF ADAPTATION**

PBA Taylor  $^{\oplus}$  has a wide range of adaptation. It is well-adapted to all the cropping regions in southern Australia.



# **PBA Taylor**<sup>(D)</sup> 'Kaspa-type' field pea

## **YIELD & ADAPTATION**

- A broad adaptation across all the cropping regions with consistently high grain yield.
- 5% to 10% higher yielding than the dual virus resistant variety PBA Wharton<sup>(b)</sup> on all soils except those high in boron and salt.
- The best variety to grow in the environments prone to virus diseases (PSbMV and BLRV).

Table 1 presents the yield of PBA Taylor<sup>()</sup> in comparison to other field pea varieties across the cropping belt in southern Australia. The yields are based on extensive field trials grown as National Variety Trials (NVT) and Pulse Breeding Australia (PBA) trials from 2017 to 2020. The yield advantage of PBA Taylor<sup>(h)</sup> has been highly consistent across cropping regions.



Table 1. Seed yield (tha) of PBA Taylor and other new pea varieties in the cropping beit of southern Australia.										u.					
Variety	Туре	NS	SW		Sou	th Austi	ralia		Vict	oria		Weste	ern Au	stralia	
		S/E	S/W	Lower EP	Mid North	South East	Upper EP	Yorke P	Mallee	Wimm- era	Ag- zone 1	Ag- zone 2	Ag- zone 3	Ag- zone 4	Ag- zone 5
PBA Taylor <sup>()</sup>	Kaspa	1.94	1.93	2.41	2.27	3.26	2.11	1.51	2.14	3.19	0.90	1.55	2.25	1.27	1.31
Kaspa <sup>⊕</sup>	Kaspa	1.53	1.62		1.67	2.75	2.06	1.43	1.99	2.84	0.93		2.08	1.14	1.40
PBA Butler	Kaspa	1.75	1.77	2.17				1.55	2.27	2.52	0.92	1.54	2.25	1.32	1.39
PBA Gunyah®	Kaspa	1.69	1.82	2.15	1.45	1.94				2.64	0.94	1.43	2.02	1.15	1.41
PBA Twilight <sup>®</sup>	Kaspa	1.67	1.81	-	1.34	-	1.58	1.44	1.81	2.66	0.94	1.41	1.99	1.13	1.40
PBA Wharton <sup>()</sup>	Kaspa	1.69	1.71	2.29	1.84	2.91	1.99	1.38	1.89	2.98	0.90	1.41	2.08		1.46
PBA Oura <sup>®</sup>	Dun				1.79	2.84	2.00	1.38	1.88	2.95	0.91	1.41	2.04	1.07	1.44
PBA Percy <sup>®</sup>	Dun	1.60		2.22	1.77	2.69	1.94	1.34	1.82	2.81	0.90	1.29	2.04	0.93	1.41
PBA Pearl®	White	1.72	1.72	2.37	1.84	3.13	2.10	1.47	2.00	3.18	0.93	1.60	2.10	1.32	1.50
Sturt	White	1.52	1.46	-	1.26	-	1.03	1.39	2.08	2.78	-	-	-	-	1.48
no. of trials		7	9	4	13	2	7	4	16	9	1	2	2	2	12

#### Table 1. Seed yield (t/ha) of PBA Taylor<sup>(b)</sup> and other field neavarieties in the cropping belt of southern</sup>

Means from reliable trials yielding on average higher than 0.5 t/ha. **Source:** National Variety Trials (NVT) and Pulse Breeding Australia (PBA) trials from 2017 to 2020.



Yield response of varieties within each region.



# **PBA Taylor**<sup>(D)</sup> 'Kaspa-type' field pea

#### AGRONOMY

PBA Taylor<sup>(h)</sup> growers should follow the same agronomic practices as for Kaspa<sup>(h)</sup>. See Table 2 for a comparison of agronomic features of current field pea varieties.

Variety		Plant vigour	Erect growth	Flowering time	Maturity time	Pod	Soil tolerance		Seed
	Plant habit	early season	habit			<b>shattering</b> at maturity	Boron	Salinity	weight (g/100)
Kaspa type									
PBA Taylor <sup>®</sup>	SD-SL	High	Fair–Good	Mid	Early–Mid	R (SP)	S	S	19
PBA Butler	SD-SL	Very High	Good	Mid–Late	Early–Mid	R (SP)	S	S	18
PBA Wharton®	SD-SL	High	Fair–Good	Early–Mid	Early	R (SP)	Т	MS	18
Kaspa <sup>⊕</sup>	SD-SL	High	Fair–Good	Late	Mid	R <i>(SP)</i>	S	S	20
PBA Gunyah <sup>®</sup>	SD-SL	High	Fair–Good	Early-Mid	Early	R (SP)	S	S	19
PBA Twilight <sup>®</sup>	SD-SL	High	Fair–Good	Early	Early	R <i>(SP)</i>	S	S	19
Australian dun t	ype								
PBA Oura <sup>®</sup>	SD-SL	High	Fair–Good	Early-Mid	Early	MR (NSP)	S	S	20
PBA Percy <sup>®</sup>	С	Very High	Poor	Early	Early	MR (NSP)	S	MT	25
Niche grain type									
PBA Pearl <sup>®</sup>	SD-SL	High	Fair–Good	Early–Mid	Early	MR (NSP)	S	MS	20
Sturt	С	Very High	Poor	Early-Mid	Mid	MR (NSP)	S	MS	-

**Key:** SD=semi-dwarf, C=conventional, SL=semi-leafless, S=susceptible, MS=moderately susceptible, MR=moderately resistant, R=resistant. SP=sugar pod type pod, NSP=non sugar pod type pod. It is important to note that seed weight varies with growing environment.

#### **DISEASE MANAGEMENT**

PBA Taylor<sup>(b)</sup> has resistance to two viruses – pea seed borne mosaic virus and bean leaf roll virus. PBA Taylor<sup>(b)</sup> is susceptible to fungal diseases (blackspot, downy mildew and powdery mildew) and bacterial blight. A proactive disease management is needed to maximise yield potential in higher risk environments.

Variety	Blackspot (Ascochyta)	Bacterial blight	<b>Downy mildew</b> (Kaspa strain)	Powdery mildew	PSbMV	BLRV (Field rating)
Kaspa type	<u>.</u>	·				
PBA Taylor <sup>®</sup>	MS	S	S	S	R	R
PBA Butler®	MS	MS	S	S	S	S
PBA Wharton <sup>()</sup>	MS	S	S	R	R	R
Kaspa®	MS	S	S	S	S	S
PBA Gunyah <sup>®</sup>	MS	S	S	S	S	S
PBA Twilight <sup>©</sup>	MS	S	S	S	S	S
Australian dun typ	e					
PBA Oura <sup>®</sup>	MS	MS	S	S	S	MR
PBA Percy <sup>®</sup>	MS	MR/MS	S	S	S	S
Niche grain type						
PBA Pearl <sup>®</sup>	MS	MS	S	S	S	R
Sturt <sup>⊕</sup>	MS	MS	S	S	S	S

Key: S=Susceptible, M=moderately, R=Resistant, PSbMV=Pea seed borne mosaic virus, BLRV=Bean leaf roll virus.




REFER TO DETAILED INFORMATION AT www.pulseaus.com.au Best management guides, crop and disease management bulletins

# **PBA Taylor** (b) (Kaspa-type' field pea


#### **GRAIN QUALITY**

PBA Taylor<sup>(b)</sup> produces medium sized spherical grain. Seed coat has a uniform tan colour similar to Kaspa<sup>(b)</sup> and is suitable for dahl and split pea production. Seed coat colour may vary slightly depending on the season.



**PBA Taylor**<sup>(D)</sup>





**Kaspa**<sup>(b)</sup>

### MARKETING

The grain is suitable to market as 'Kaspa-type' grain, which is exported to the Asian sub-continent for production of dahl, flour and roasted snack foods.

The grain is also suitable for stockfeed.

# BREEDING

PBA Taylor<sup>()</sup> (tested as OZP1408) was identified by PBA field pea team and is derived from two breeding lines. The crossing was made in 2006 at Department of Jobs, Precincts and Regions (DJPR), Horsham. Several cycles of selections were made to improve yield stability and disease resistance. PBA Taylor<sup>()</sup> is named after Taylors Beach in south west Victoria.

# **PULSE AGRONOMY**

Agronomy and disease management information has been developed with the assistance of 'Southern region pulse agronomy project' co-funded by GRDC, SARDI, DJPR and NSW-DPI.

# FIELD PEA BREEDING AUSTRALIA

Pulse Breeding Australia (PBA) was an unincorporated joint venture between the GRDC, University of Adelaide, University of Sydney, SARDI, Agriculture Victoria Research, NSW DPI, DAF (QLD), DPIRD WA and Pulse Australia.

#### FOR MORE INFORMATION

**GRDC** PO Box 536 Kingston ACT 2604 Ph: 02 6166 4500 Field Pea Breeder Dr Babu Pandey Agriculture Victoria 110 Natimuk Road Horsham, VIC 3400

Ph: 03 4344 3157 babu.pandey@agriculture.vic. gov.au

Seednet

Planting Productivity

Ph: 1300 799 246

Fax: 03 5381 0490

#### **SEED ENQUIRIES**

Seednet National Production and Logistics Office 7 Golf Course Rd PO Box 1409, Horsham Vic 3402

#### Eastern Australia

Stuart Ockerby Sth NSW, VIC, TAS, SA Ph: 0448 469 745 stuart.ockerby@seednet.com.au

Seednet's mission is:

"To deliver high performance seed based genetics to Australian grain growers and end user customers via superior product and service delivery channels".

Seednet is proud to partner with Field Pea Breeding Australia and invest in the improvement of Australian field pea varieties.

#### **AGRONOMIC ENQUIRIES**

South Australia Penny Roberts, SARDI, SA Ph: 08 8841 2401 Penny.Roberts@sa.gov.au

#### Victoria

Jason Brand, Agriculture Victoria, Horsham, Vic Ph: 03 5362 2341 Jason.Brand@agriculture.vic.gov.au

#### Southern New South Wales

Mark Richards, NSW Department of Primary Industries, Wagga Wagga, NSW Ph: 0428 630 429 mark.richards@dpi.nsw.gov.au

Disclaimer: Recommendations have been made from information available to date and considered reliable, and will be updated as further information comes to hand. Readers who act on this information do so at their own risk. No liability or responsibility is accepted for any actions or outcomes arising from use of the material contained in this publication. Reproduction of this brochure in any edited form must be approved by the National Field Pea Breeding Program © 2021

Version September/2021

admin@seednet.com.au www.seednet.com.au

Western Australia David Clegg Ph: 0408 630 641 david.clegg@seednet.com.au